

A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.

Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.

CONF

NOTICE

PORTIONS OF THIS REPORT ARE ILLLEGIBLE. It has been reproduced from the best available copy to permit the broadest possible availability.

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36.

CONF-840842--3

TITLE: MAXIMUM SUPERCOOLING IN LIQUID ^3He - ^4He MIXTURES NEAR THE TRICRITICAL POINT

LA-UR--84-904

AUTHOR(S): Dipen N. Sinha and James K. Hoffer

DE84 010076

SUBMITTED TO: 17th International Conference on Low Temperature Physics
Karlsruhe, West Germany, August 1984.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

MAXIMUM SUPERCOOLING IN LIQUID ^3He - ^4He MIXTURES NEAR THE TRICRITICAL POINT*

Dipen N. SINHA

Rockwell International, Anaheim, CA 92803, U.S.A.

James K. HOFFER

Los Alamos National Laboratory, Los Alamos, NM 87545, U.S.A.

Measurements of supercooling in liquid ^3He - ^4He mixtures near the tricritical point are presented. The reduced temperature range $0.001 < \epsilon \equiv (1 - T/T_c) < 0.01$ was investigated for three different rates of cooling using a pressure-quench technique. For $\epsilon < 0.012$, the maximum supercooling was found to be a function of the cooling rate. Comparisons with data in organic binary mixtures are given.

1. INTRODUCTION

We have made quantitative measurements of the maximum supercooling δT_m in the superfluid phase of liquid ^3He - ^4He mixtures at three different quench rates in the vicinity of the tricritical point T_c . A pressure quench technique (1) was used to produce supercooling in the mixture and the sudden attenuation in the intensity of a transmitted laser beam through the mixture was used to determine the cloud-point, which marks the limit of supercooling δT_m . Employing a similar technique, Alpern et al. (2) recently found that as the tricritical point is approached, i.e., as $\epsilon \equiv (1 - T/T_c) \rightarrow 0$, the normalized maximum supercooling $\delta T_m/\Delta T$ tends to diverge. We use the definitions $\delta T_m \equiv T_{ps} - T_{cp}$ and $\Delta T \equiv T_c - T_{ps}$, where T_{ps} is the equilibrium phase separation temperature and T_{cp} is the measured cloud-point temperature. Such behavior of supercooling near the critical point has been observed for organic binary mixtures (3), but Alpern et al. showed the divergence of $\delta T_m/\Delta T$ to be much stronger in the case of ^3He - ^4He . They also pointed out that this feature of stronger divergence of the supercooling near T_c could be a peculiarity of tricritical systems. In this paper, we show that the degree of divergence of $\delta T_m/\Delta T$ as $\epsilon \rightarrow 0$ is a function of the quench rate, i.e., the rate of cooling; and if one compares the ^3He - ^4He data with those of organic mixtures for equivalent quench rates, then no meaningful discrepancy is observed. We also show other similarities in the results between ^3He - ^4He and other studies in organic binary mixtures (4).

The divergence of supercooling as $\epsilon \rightarrow 0$ can be explained in terms of critical slowing-down of diffusion processes (5,6). The experimentally determined cloud-point indicates the stage of

phase-separation when the evolving minority-phase clusters or droplets have grown large enough to begin to scatter light strongly. Consequently, the cloud-point phenomena always lag in time behind the actual onset of homogeneous nucleation. Because of critical slowing down, this time-lag gets progressively larger as one approaches the critical point. Coupled with the finite quench rates used in actual experiments, this appears to indicate additional supercooling.

2. EXPERIMENTAL

The supercooling of the liquid ^3He - ^4He mixture in the metastable region of the miscibility gap below the tricritical point was produced using a pressure quench technique (1). Pressurization of the ^3He - ^4He mixture inside the sample cell was accomplished by pure ^4He actuating two horizontal bellows moving toward each other. A germanium resistance thermometer monitored the temperature T and an in-situ capacitive transducer with a resolution of 10^{-5} atm monitored the pressure P of the mixture. The intensity I_0 of the transmitted He-Ne laser beam, attenuated to approximately 0.1 mW before entering the cell, was measured using a sensitive photodiode. Different pressure quench rates were made possible by using several parallel and independently selectable fixed impedances in the pressure release line of the pressurization bellows outside the cryostat. Fig. 1 shows a typical pressure quench history. The cloud-point is indicated by the point where I_0 suddenly starts to decrease. The actual supercooling δT_m was evaluated from the pressure at the cloud-point and the well-known pressure dependence of the tricritical point (1,2).

*Work supported in part by the U. S. Department of Energy.




Figure 1. Time dependence of the pressure, temperature, and forward transmission intensity during a pressure-induced quench into the miscibility gap of a ^3He - ^4He liquid mixture.

3. RESULTS AND DISCUSSION

We now discuss the results of our experiment to determine the homogeneous nucleation limit to supercooling. Three different rates of supercooling were used: 67, 27, and 12 mK/s. These are the average values of the cooling rates, determined from the pressure quench data, at the moment the light intensity I_0 begins to drop (see Fig. 1). All the measurements were made in the superfluid phase of the mixture. Fig. 2 shows the reduced supercooling $\delta T_m/\Delta T$ as a function of ϵ on a log-log plot for the three different quench rates. The long-dashed line is the data of Ref. 4 for a quench rate of 30 mK/s in a lutidine-water mixture. The short-dashed line represents the data of Ref. 3 for lutidine-water using effective quench rates of less than 1 mK/s. The dotted line represents data in the ^4He phase of ^3He - ^4He mixtures from Ref. 2 for quench rates greater than 150 mK/s. For values of $\epsilon < 0.012$, the dependence on quench rate is obvious. All data can be fairly well represented by straight lines. At $\epsilon > 0.012$, the quench rate dependence of the supercooling gradually diminishes and the $\delta T_m/\Delta T$ data become consistent with the prediction from classical nucleation theory (dot-dashed line). Similar behavior was observed by Strey et al. (4) in lutidine-water for three different quench rates used in their experiments (see Fig. 8 of Ref. 4). Considering the differing experimental techniques used (pressure quench vs. temperature quench) and the disparate thermodynamic properties of the two systems, no meaningful inconsistency can be inferred from data in organic mixtures in comparison to either our data or those of Benda et al. in ^3He - ^4He , i.e., there is no unreasonably stronger divergence in the ^3He - ^4He supercooling data as $\epsilon \rightarrow 0$. However, the effects of the particular experimental techniques used,




Figure 2. Relative supercooling vs reduced temperature for three different cooling rates in the superfluid phase of a ^3He - ^4He mixture of tricritical composition.

such as pressure quench vs temperature quench, in determining the cloud-point, have not been adequately studied. Therefore, we do not feel that there is sufficient evidence to suggest that the divergence of supercooling near the tricritical point in ^3He - ^4He is substantially different from that in organic mixtures near their critical points. In conclusion, we have shown that in ^3He - ^4He mixtures, the anomalous maximum supercooling is largely an artifact of the finite rates of cooling inherent to the experimental methods that have been utilized.

REFERENCES

- (1) D. N. Sinha and J. K. Hoffer, Phys. Rev. Lett. 50 (1983) 515.
- (2) P. Alpern, Th. Benda, and P. Leiderer, Phys. Rev. Lett. 49 (1982) 1267.
- (3) A. J. Schwartz, S. Krishnamurthy, and W. I. Goldburg, Phys. Rev. 21A (1980) 1331.
- (4) R. Strey, J. Wagner, and J. Woerman, Ber. Bunsenges. Phys. Chem. 86 (1982) 306.
- (5) K. Binder and D. Stauffer, Adv. Phys. 25 (1976) 343.
- (6) J. S. Langer and A. J. Schwartz, Phys. Rev. 21A (1980) 948.